

## Case Report

# Unsuspected Postpartum Complication: A Case Report of a Post Cesarean Gastric Rupture

Carolyn M Zahler-Miller, ND<sup>1</sup> , Rasmita Jalla, MD<sup>2</sup>, Louise Wannamaker, MD<sup>1</sup>

<sup>1</sup> Obstetrics and Gynecology, Augusta University, <sup>2</sup> Internal Medicine, Augusta University

Keywords: intellectual impairment, gastric perforation, gastric necrosis, gastric dilatation, diabetes mellitus, cesarean delivery

<https://doi.org/10.54053/001c.121487>

---

## North American Proceedings in Gynecology & Obstetrics

---

**Background:** Although rare, acute gastric dilatation is associated with life threatening complications. While several cases have been documented in various settings, there are very few cases documented in association with gynecological surgeries.

**Objective:** We report a case of acute gastric dilatation with subsequent necrosis and perforation in 35-year-old postpartum patient two days following a primary cesarean section. The patient's medical history was significant for diabetes and intellectual disability. Despite treatment and resuscitative measures following gastric rupture, the patient continued to decompensate, suffering respiratory failure, anoxic brain injury, and subsequently death.

**Methods:** Case Report

**Results:** n/a

**Conclusion:** Gastric necrosis is a postoperative complication any surgeon should be aware of. While there are very few documented cases of gastric necrosis and perforation after a gynecological surgery, pelvic surgeons should still be cognizant of this severely fatal complication. Obstetricians must maintain a high suspicion of ileus in any post-cesarean patient with nausea or vomiting, particularly in patients with a history of diabetes mellitus and intellectual impairment who are at risk for gastric necrosis.

Awareness of this potential complication may help providers initiate treatment modalities quickly, thus reducing mortality.

## CASE

A 35-year-old African American woman G1P0 diagnosed with intellectual impairment, pre-eclampsia and type II diabetes mellitus initially presented to Labor and Delivery for a scheduled induction of labor at 37 weeks 3 days of gestation. She had a Bishop score of 7 and was started on oxytocin on December 26, 2018. 24 hours into her induction, the fetal heart tracing became non-reassuring and the decision was made to proceed with a cesarean section. She underwent a primary low transverse cesarean section and bilateral tubal ligation without any complications, delivering a singleton at 37 weeks 4 days. Her peripartum status was stable for the next day and a half.

Early on the second morning after delivery, the patient developed diarrhea and vomiting with approximately 200 mL of non-bilious emesis. By 2350 that day, her oxygen sat-

uration decreased to 90%. She became progressively more uncomfortable with mild tachypnea and increasing shortness of breath. By 0030, she was also found to be diaphoretic and tachycardic. Her EKG at that time showed a prolonged QT interval with a rate in the 120s. It was suspected she had a pulmonary embolus, so she was anticoagulated per protocol and placed on 2 liters of oxygen via nasal cannula while awaiting a CT Angiogram.

As the patient was being prepped for a CT scan, she was noted to have a rapid change in status with acute onset abdominal distension, elevation of heart rate to the 150s, and worsening tachypnea. She additionally had visible retractions with respirations and tympany on abdominal exam. She was transferred to the Intensive Care Unit (ICU) and intubated. At this point, the patient was continued on low molecular weight heparin for anticoagulation. The CT scan could not be obtained due to her unstable condition. The

patient then suffered cardiac arrest. Return of spontaneous circulation was achieved with cardiopulmonary resuscitation. A chest x-ray showed free air under the diaphragm and a severely enlarged stomach. Acute care surgery was consulted and decision was made to proceed with emergent exploratory laparotomy in the ICU as the patient was not stable enough to transfer to the operating room.

Upon entering the abdomen, the patient was found to have acute gastric necrosis thought to be secondary to acute gastric dilatation. Three liters of turbid fluid with food particles were evacuated. The small bowel was ischemic and dusky throughout the entire length and grossly dilated with no perforation or injury. The colon appeared normal and intact. Two-thirds of the anterior and proximal stomach was necrotic with a nasogastric tube perforating through a large area of necrosis. A subtotal gastrectomy followed. The spleen showed multiple areas of bleeding, so it was removed as well to prevent any further complications. After this, a washout of the abdomen was done and a temporary closure device was placed with plans to return 24-48 hours later for a second look after the patient was stabilized and resuscitated. The patient had immediate improvement in oxygenation, but remained critically ill on maximum doses of three pressors. She was subsequently transferred to the surgical ICU.

At 0919 on postpartum day 5 (post-operative day 2 from the exploratory laparotomy), the patient remained intubated off sedation. All reflexes were absent and she did not withdraw to pain. The family decided to withdraw care, and she was extubated and taken off pressors at 0930. The patient passed away at 0937. Cause of death was determined to be ileus and severe gastric dilatation leading to abdominal compartment syndrome with acute gastric necrosis and anoxic brain injury leading to respiratory failure.

## DISCUSSION

Acute gastric dilatation leading to subsequent necrosis and perforation is an exceedingly rare complication of abdominal surgery. While there have been several previous reports of gastric necrosis or perforation in pregnancy related to ascariasis or other unknown causes, the literature has no documented cases of gastric perforation in post-partum patients with diabetes. The stomach has an abundant blood supply from many vessels and collateral circulation, so it is rare for gastric necrosis to occur. However, there are risk factors such as gastric outlet obstruction, retroperitoneal tumors, anorexia nervosa and bulimia, psychogenic reasons, diabetes mellitus, trauma, electrolyte disorders, cerebral palsy, diaphragmatic hernia, and gastric volvulus that can lead to its occurrence (Powell et al. 2003). The likely sequence of events was that our patient developed a severe ileus and subsequent gastric dilatation leading to abdominal compartment syndrome, and that her history of di-

abetes and potential impaired vascularity of the stomach wall placed her at greater risk for gastric necrosis. Thus, she sustained a gastric perforation upon placement of the NG tube in the ICU setting.

In regards to this patient's intellectual impairment, it may have been a potential risk factor as well. Psychogenic and neurologic factors can influence behaviors, which may lead to severe complications such as gastric necrosis (Powell et al. 2019). Examples of a psychogenic factors are pathologic aerophagia, which is swallowing excessive amounts of air, and overeating due to impaired intrinsic mechanisms which alert the person is full (Powell et al. 2019; Zvizdic et al. 2019). It also is possible that our patient was unable to adequately verbalize symptoms such as abdominal pain or difficulty breathing due to her mental disability.

Signs and symptoms of gastric necrosis include vomiting, abdominal pain, abdominal bloating, and distension. Imaging may reveal a severely enlarged stomach with free air under the diaphragm, which suggests perforation. If a diagnosis of gastric necrosis or perforation is made, swift actions must be taken due to the high mortality rate of 50-80% (Zvizdic et al. 2019). Treatment is emergent surgery with the procedure usually including surgical decompression, partial gastrectomy or total gastrectomy and esophago-jejunostomy (Gündeş et al. 2013). Delayed treatment may lead to worse prognosis, so it is imperative to initiate treatment as soon as possible. Many patients die even with efficient treatment due to multisystem organ failure and sepsis (Ntirushwa et al. 2016; Powell et al. 2019). Unfortunately, our patient became hypoxemic secondary to respiratory failure from her abdominal compartment syndrome, which led to circulatory collapse requiring cardiopulmonary resuscitation. Due to her prolonged hypoxemia, she suffered anoxic brain injury and ultimately brain death.

## CONCLUSION

Although rare, gastric necrosis and perforation are life threatening postoperative conditions if they occur. Quick treatment modalities must be implemented in order to reduce mortality. Obstetricians must maintain a high suspicion of ileus in any post-cesarean patient with nausea or vomiting, particularly in patients with a history of diabetes mellitus and intellectual impairment who are at risk for gastric necrosis (Ntirushwa et al. 2016; Powell et al. 2019). Early imaging is crucial. If a nasogastric tube needs to be placed, care needs to be taken to avoid perforation.

Submitted: June 20, 2021 EDT, Accepted: June 26, 2021 EDT



This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CCBY-4.0). View this license's legal deed at <http://creativecommons.org/licenses/by/4.0> and legal code at <http://creativecommons.org/licenses/by/4.0/legalcode> for more information.

## REFERENCES

- Gündeş, E., T. Küçükkartallar, A. Tekin, and M. Çakır. 2013. "Gastric Necrosis and Perforation Caused by Acute Gastric Dilatation." *Ulus Cerrahi Derg* 30 (3): 179–81. <https://doi.org/10.5152/UCD.2013.34>.
- Ntirushwa, D., S. Rulisa, F. Muhorakeye, L. Bazzett-Matabele, T. Rurangwa, and M. Small. 2016. "Gastric Rupture in Pregnancy: Case Series from a Tertiary Institution in Rwanda and Review of the Literature." *AJP Rep* 6 (4): e436–41. <https://doi.org/10.1055/s-0036-1597619>.
- Powell, John L., Joseph Payne, Clinton L. Meyer, Paul R. Moncla, Chikako Murakami, Wataru Irie, Chizuko Sasaki, et al. 2019. "Extensive Gastric Necrosis Secondary to Acute Gastric Dilatation: A Case Report." *Legal Medicine* 36:85–88. <https://doi.org/10.1016/j.legalmed.2018.11.007>.
- Powell, John L., Joseph Payne, Clinton L. Meyer, and Paul R. Moncla. 2003. "Gastric Necrosis Associated with Acute Gastric Dilatation and Small Bowel Obstruction: Case Report." *Gynecologic Oncology* 90 (1): 200–203. [https://doi.org/10.1016/S0090-8258\(03\)00204-X](https://doi.org/10.1016/S0090-8258(03)00204-X).
- Zvizdic, Zlatan, Asmir Jonuzi, Aleksandra Djuran, and Semir Vranic. 2019. "Gastric Necrosis and Perforation Following Massive Gastric Dilatation in an Adolescent Girl: A Rare Cause of Acute Abdomen." *Frontiers in Surgery* 6:3. <https://doi.org/10.3389/fsurg.2019.00003>.